

Written by Guy Mong Ky TRAN

Saturday, 04 November 2017 17:31 - Last Updated Saturday, 04 November 2017 17:33

There are no translations available.

Highly pathogenic Influenza virus A H5N1 Indonesia & Vietnam Hemagglutinin (HA1) contains an urokinase-plasminogen activator explaining the fibrinolysis

Présentation Poster au [8TH EUROPEAN MEETING ON VIRAL ZOOSES\(Saint Raphaël, France, Octobre 2017\)](#)

Guy Mong Ky Tran(1) & Adrien Caprani(2)

(1) Retired since 2014 from University of Auvergne Rhone-Alpes, Health Regional Agency, Public Health Department, Hospital Hotel-Dieu, Clermont-Ferrand ; 31 Av du Bois - Chatenay Malabry, France. email: mkg_tran@yahoo.fr

(2) Association "Positifs", 147 chemin de la futaie, 83550 Vidauban, email: positifpresident@yahoo.fr

Avian Influenza is compared to a chicken lethal Ebola for its internal hemorrhagic symptoms. Oseltamivir is inefficient and not recommended in hemorrhagic influenza. In 1996, Zhilinskaia IN (Vopr Virusol) found by computer analysis a similarity between Influenza virus hemagglutinin HA and plasminogen activator (PA), explaining the fibrinolysis. We analysed further these results in Influenza virus A H5N1 Indonesia (83% mortality) and Vietnam HA1. We compared to urokinase (U-PA), tissue PA, TSV-PA, Batroxobin. Results : All the 3 active site residues of serine protease (His, Asp, Ser) were found and the best match was with U-PA :

C.113-Highly pathogenic Influenza virus A H5N1 Indonesia & Vietnam Hemagglutinin (HA1) contains an u

Written by Guy Mong Ky TRAN

Saturday, 04 November 2017 17:31 - Last Updated Saturday, 04 November 2017 17:33

H5N1 Indonesia/Vietnam IPKSS-W--S-SH

U-PA/t-PA active site His ILISScWviSaTH

H5N1 Indonesia/Vietnam G I HHpNDAAEQTK

U-PA/t-PA active site Asp (A,L)HH NDI ALQ IR

H5N1 Vietnam KGDS- - TIMKS(L,E)EYGNCKCQTPMGA I

U-PA/t-PA active site Ser QGDSggPLVcS L Q WIRSHTKGEE- NG(A,L)

H5N1 Indonesia/Vietnam 141-YLGKSSF(R,F)N-150

U-PA YLGR-S(L,R)L N

Conversely, Influenza H7 China HA1 has His replaced by Arg, Asp replaced by Ser. The motif His-Phe (HF), in the pocket binding Plasminogen clivage site Arg-Ala-Arg, was present in Influenza H5N1 and U-PA (319-HF-320); but not in Influenza virus H7 China, replaced by Gly-Gly (inactive). Conclusion : The plasminogen activator in avian Influenza virus H5N1 hemagglutinin HA1 (Zhilinskaia) was confirmed and may explain fibrinolysis in hemorrhagic Influenza. It seems rationale to use plasminogen inhibitors to block this fibrinolysis. A peptidomimetic drug of the plasminogen Arg-Ala-Arg (RAR) at the cleavage site Arg / Ala can be designed by docking. This strategy was successful in HIV-1 tritherapy, with FP peptidomimetics.

C.113-Highly pathogenic Influenza virus A H5N1 Indonesia & Vietnam Hemagglutinin (HA1) contains an u

Written by Guy Mong Ky TRAN

Saturday, 04 November 2017 17:31 - Last Updated Saturday, 04 November 2017 17:33
